Lecture 5. Epigenomics and regulation of genome activity

Dep. Biotechnology Prof. Kenzhebayeva S.S.

Aim of the Lecture. Expected Learning Outcomes

Aim of the Lecture

To introduce the fundamental principles of epigenomics and the molecular mechanisms regulating genome activity, with emphasis on DNA methylation, histone modifications, chromatin organization, modern epigenomic technologies, and their applications in functional genomics and biotechnology.

Expected Learning Outcomes

- Define epigenetics and epigenomics and explain their significance in postgenomic biology.
- Describe the main mechanisms of epigenetic regulation, including DNA methylation, histone modifications, and chromatin remodeling.
- Explain the roles of chromatin organization and accessibility in regulating gene expression.
- Distinguish between genetic and epigenetic changes.
- Describe the principles of modern epigenomic technologies, including bisulfite sequencing, ChIP-seq, and ATAC-seq.
- Explain the relationships between epigenetic regulation, plant development, and environmental stress responses.
- Discuss the potential inheritance and reversibility of epigenetic modifications.
- Identify applications of epigenomics in plant, medical, and industrial biotechnology.
- Evaluate the opportunities and limitations of epigenome editing and multi-omics integration.
    - ***Key learning outcome: Understanding how epigenetic mechanisms regulate genome activity and contribute to biological development, adaptation, and phenotypic diversity.***

Lecture Plan

1. Introduction to Epigenetics and Epigenomics: Definitions, historical development, and biological significance.
2. Chromatin Structure and Genome Organization: Nucleosomes, chromatin states, and genome accessibility.
3. DNA Methylation and Demethylation: Molecular mechanisms and roles in gene regulation.
4. Histone Modifications and Chromatin Remodeling: Epigenetic marks and regulation of transcription.
5. Non-Coding RNAs and Epigenetic Regulation: Small RNAs, RNA-directed DNA methylation, and regulatory interactions.
6. Modern Epigenomic Technologies: Bisulfite sequencing, ChIP-seq, ATAC-seq, and epigenomic data analysis.
7. Applications in Biotechnology: Plant development, stress responses, epigenetic variation, and crop improvement.
8. Challenges and Future Perspectives: Epigenome editing, epigenetic inheritance, and multi-omics integration.

Introduction to Epigenetics and Epigenomics

Epigenetics is the study of mechanisms that regulate gene activity and cellular states without changing the underlying DNA nucleotide sequence.

Epigenomics investigates epigenetic modifications and chromatin organization across the entire genome.

Main components of epigenetic regulation:

- DNA methylation: Chemical modification of DNA bases, primarily cytosine.
- Histone modifications: Acetylation, methylation, phosphorylation, and other modifications of histone proteins.
- Chromatin remodeling: Changes in nucleosome positioning and DNA accessibility.
- Regulatory RNAs: Non-coding RNAs involved in chromatin-associated gene regulation.

Major objectives:

- Understanding genome-wide regulation of gene expression.
- Investigating cellular differentiation and developmental processes.
- Studying environmental influences on genome activity.
- Identifying epigenetic mechanisms associated with phenotypic variation.

Key concept: Epigenomics explains how cells with similar DNA sequences can exhibit different gene expression patterns and biological functions.

Genetic vs. Epigenetic Regulation

Genetic and epigenetic mechanisms contribute to biological variation through different molecular processes.

Characteristic	Genetic regulation	Epigenetic regulation
Molecular basis	DNA sequence variation	DNA and chromatin modifications
DNA sequence	May be altered by mutations	Generally unchanged
Main mechanisms	Mutations, allelic variation, regulatory sequence differences	DNA methylation, histone modifications, chromatin remodeling
Stability	DNA variants are usually stable	Epigenetic states may be stable or dynamic
Inheritance	Transmitted through DNA replication and reproduction	Some states can persist through cell division or generations

Example in plant biotechnology:

- Genetic variation: A mutation in a stress-responsive gene may alter its encoded protein or regulatory sequence.
- Epigenetic variation: Changes in DNA methylation near a stress-responsive gene may influence its transcription without changing the DNA sequence.

Key concept: Genetic variation changes DNA sequence, whereas epigenetic regulation modifies how genomic information is accessed and expressed.

Biological Significance of Epigenetic Mechanisms

Epigenetic mechanisms are essential for coordinating gene activity during development and environmental adaptation.

Major biological functions:

- Cell differentiation: Establishment and maintenance of tissue-specific gene expression patterns.
- Plant development: Regulation of germination, flowering, organ formation, and reproductive development.
- Genome stability: Silencing of transposable elements and repetitive DNA sequences.
- Environmental responses: Modulation of gene expression under drought, salinity, temperature extremes, and pathogen infection.
- Developmental plasticity: Adjustment of gene activity in response to internal and external signals.
- Epigenetic memory: Persistence of selected regulatory states after an environmental stimulus, under certain conditions.

Applications in biotechnology:

- Identification of epigenetic markers associated with important traits.
- Investigation of stress-responsive regulatory pathways.
- Development of approaches for manipulating gene activity without modifying DNA sequence.

Key concept: Epigenetic regulation contributes to genome stability, developmental coordination, and phenotypic plasticity.

Chromatin Structure and Nucleosome Organization



Chromatin is a complex of DNA, histone proteins, and associated molecules that organizes the genome within the nucleus.

Basic structural unit — nucleosome:

Approximately 147 base pairs of DNA are wrapped around a histone octamer.

The histone octamer contains two copies each of H2A, H2B, H3, and H4.

Linker DNA connects neighboring nucleosomes.

Histone H1 contributes to chromatin organization and compaction.



Levels of chromatin organization:

1. DNA double helix.
2. Nucleosomes arranged along DNA.
3. Higher-order chromatin organization and interactions.
4. Chromosome territories and three-dimensional genome architecture.

Functional significance:

- DNA packaging and protection.
- Regulation of transcription factor accessibility.
- Coordination of DNA replication and repair.
- Control of gene expression through chromatin modification

Key concept: Nucleosome positioning and chromatin organization influence the accessibility of DNA to transcriptional regulators and other molecular machinery.

Euchromatin, Heterochromatin, and Chromatin Accessibility

Chromatin accessibility describes the extent to which DNA is available for interaction with transcription factors and other regulatory proteins.

Major chromatin states:

- Euchromatin: Generally less condensed, gene-rich, and associated with transcriptionally permissive regions.
- Heterochromatin: More compact, often enriched in repetitive DNA and transposable elements, and frequently associated with transcriptional silencing.
- Facultative heterochromatin: Chromatin that can transition between repressed and more active states depending on cell type or developmental conditions.
- Constitutive heterochromatin: Relatively stable compact chromatin, commonly found near centromeres and repetitive regions.

Factors influencing chromatin accessibility:

- Nucleosome positioning and occupancy.
- Histone modifications and histone variants.
- DNA methylation.
- ATP-dependent chromatin remodeling complexes.

Key concept: Chromatin accessibility influences the ability of regulatory proteins to interact with DNA and control transcription.





DNA Methylation: Principles and Biological Functions

DNA methylation is an epigenetic modification involving the addition of a methyl group to DNA bases, predominantly the fifth carbon of cytosine, forming 5-methylcytosine (5mC).

Major characteristics:

- Catalyzed by DNA methyltransferase enzymes.
- Occurs in different sequence contexts depending on the organism.
- Can influence gene expression, chromatin organization, and genome stability.
- Often contributes to transcriptional silencing of transposable elements and repetitive sequences.
- May be maintained during DNA replication or dynamically modified during development and environmental responses.

• Biological functions:

- Regulation of gene expression.
- Suppression of transposable element activity.
- Maintenance of genome integrity.
- Control of developmental and tissue-specific processes.
- Participation in responses to environmental stimuli.

Important distinction: DNA methylation is not universally

Key concept: DNA methylation is an important mechanism for regulating genome activity and maintaining epigenetic states.





DNA Methylation Patterns in Plants

Plant genomes exhibit DNA methylation in three major cytosine sequence contexts.

Methylation context	Sequence	Main maintenance mechanism
CG	Cytosine followed by guanine	MET1
CHG	H = A, T, or C	CMT3, associated with H3K9 methylation
CHH	H = A, T, or C	DRM2/RdDM and CMT2-dependent pathways

Biological significance:

- Transposable element silencing.
- Regulation of gene activity.
- Maintenance of genome stability.
- Developmental and environmental responses.

Key concept: Plant DNA methylation is distinguished by its occurrence in CG, CHG, and CHH contexts and by specialized maintenance pathways.

Major characteristics of plant DNA methylation:

- CG methylation is common in transposable elements and some gene bodies.
- CHG and CHH methylation are frequently enriched in transposable elements and repetitive sequences.
- RNA-directed DNA methylation (RdDM) contributes to the establishment of methylation, particularly in CHH contexts.
- DNA methylation patterns can differ between tissues, developmental stages, and environmental conditions.

DNA Demethylation and Epigenetic Dynamics

DNA demethylation is the removal or loss of DNA methylation marks, allowing epigenetic states to change during development and environmental responses.

Main mechanisms:

1. Active demethylation: Enzymatic removal of methylated cytosines through DNA glycosylase-initiated base excision repair.
2. Passive demethylation: Progressive loss of methylation when maintenance methylation is insufficient during DNA replication.

- Major plant DNA demethylation enzymes:
- ROS1 (REPRESSOR OF SILENCING 1).
- DME (DEMETER).
- DML2 and DML3 (DEMETER-LIKE proteins).

Biological functions:

- Regulation of developmental gene expression.
- Maintenance of appropriate methylation patterns.
- Control of selected transposable elements and nearby genes.
- Contribution to reproductive development.
- Participation in responses to environmental changes.

Histone Modifications: Major Types and Functions

Histone modifications are chemical changes to histone proteins that influence chromatin structure, protein interactions, and genome activity.

Major types of histone modifications:

- **Acetylation:** *Addition of acetyl groups to lysine residues, commonly associated with transcriptionally active chromatin.*
- **Methylation:** *Addition of methyl groups to lysine or arginine residues; effects depend on the modified residue and methylation state.*
- **Phosphorylation:** *Addition of phosphate groups, involved in DNA damage responses, chromosome condensation, and signaling.*
- **Ubiquitination:** *Attachment of ubiquitin to histones, contributing to transcriptional regulation and DNA repair.*
- **SUMOylation:** *Attachment of SUMO proteins, influencing chromatin-associated regulatory processes.*

Major regulatory proteins:

- Writers: Enzymes that add histone modifications.
- Erasers: Enzymes that remove modifications.
- Readers: Proteins that recognize specific histone marks.

Functional significance:

- Regulation of chromatin accessibility.
- Recruitment of transcriptional activators and repressors.
- Control of developmental and stress-responsive gene expression.
- Maintenance of chromatin states and genome stability.

Key concept: Histone modifications influence gene expression through their effects on chromatin structure and interactions with regulatory proteins.

Biological significance: Histone modifications participate in gene activation, transcriptional repression, DNA repair, and developmental regulation



DNA Demethylation and Epigenetic Dynamics

NA Methylation Dynamics

Methylation

Addition or maintenance of methylation marks
(MET1, CMTs, DRM2)

Demethylation

Removal or loss of methylation marks
(ROS1, DME, DMLs)

Dynamic regulation of DNA methylation patterns and genome activity

Key concept: The balance between DNA methylation and demethylation contributes to the establishment, maintenance, and modification of epigenetic states.

Histone Acetylation and Methylation in Gene Regulation

Histone acetylation and methylation are two major epigenetic mechanisms involved in the regulation of transcription.

Important histone marks:

- *H3K4me3: Frequently associated with transcriptionally active genes.*
- *H3K27me3: Associated with Polycomb-mediated gene repression.*
- *H3K9me2: Commonly associated with heterochromatin and transposable element silencing in plants.*
- *H3K9ac: Often enriched in transcriptionally active chromatin.*

Applications in plant biotechnology:

- Investigation of developmental gene regulation.
- Characterization of chromatin responses to environmental stress.
- Identification of epigenetic changes associated with phenotypic variation.

Key concept: Histone acetylation generally favors transcriptional activity, whereas histone methylation can be associated with either activation or repression.

Characteristic	Histone acetylation	Histone methylation
Chemical modification	Addition of acetyl groups	Addition of methyl groups
Major enzymes	Histone acetyltransferases (HATs)	Histone methyltransferases (HMTs)
Removal	Histone deacetylases (HDACs)	Histone demethylases (HDMs)
Typical effects	Often associated with active transcription	Activation or repression depending on the mark
Examples	H3K9ac, H3K27ac	H3K4me3, H3K27me3, H3K9me2

Relationships between histone marks and transcription

Active chromatin

H3K9ac, H3K4me3

Often associated with transcriptionally active genes

Repressive chromatin

H3K27me3, H3K9me2

Often associated with transcriptional repression

These are common associations, not universal rules; the effect of a histone mark depends on genomic and cellular context.

ATP-Dependent Chromatin Remodeling

ATP-dependent chromatin remodeling is a molecular process in which specialized protein complexes use energy from ATP hydrolysis to alter nucleosome positioning, composition, or organization.

Major remodeling mechanisms:

- Nucleosome sliding: Repositioning nucleosomes along DNA.
- Nucleosome eviction: Removal of nucleosomes to increase DNA accessibility.
- Histone exchange: Replacement of canonical histones with histone variants.
- Chromatin reorganization: Modification of local nucleosome arrangements.
- Major chromatin remodeling families:
- SWI/SNF: Involved in nucleosome repositioning and transcriptional regulation.
- ISWI: Contributes to nucleosome spacing and chromatin organization.
- CHD: Participates in transcriptional regulation and chromatin organization.
- INO80: Involved in histone variant exchange, DNA repair, and genome stability.

Biological functions:

- Regulation of transcription factor accessibility.
- Control of gene expression during development.
- Coordination of DNA replication and repair.
- Modulation of plant responses to environmental stress.



Non-Coding RNAs in Epigenetic Regulation

Non-coding RNAs (ncRNAs) are RNA molecules that are not translated into proteins but participate in the regulation of gene expression, chromatin organization, and genome stability.

Major types of regulatory non-coding RNAs:

- ❑ MicroRNAs (miRNAs): Small RNAs, typically 20–24 nucleotides in plants, that regulate gene expression mainly through target mRNA cleavage or translational repression.
- ❑ Small interfering RNAs (siRNAs): Small RNAs involved in RNA silencing, transposable element suppression, and epigenetic regulation.
- ❑ Long non-coding RNAs (lncRNAs): Transcripts generally longer than 200 nucleotides that can participate in transcriptional and chromatin-associated regulation.
- ❑ Natural antisense transcripts (NATs): RNAs transcribed from the opposite strand of overlapping genomic regions that may influence transcript abundance and regulation.

Biological functions:

- Regulation of developmental and stress-responsive genes.
- Control of transposable element activity.
- Recruitment or guidance of chromatin-modifying complexes in specific pathways.
- Maintenance of genome stability.
- Important distinction: Not all non-coding RNAs regulate chromatin directly; many function primarily through post-transcriptional mechanisms.

Key concept: Regulatory non-coding RNAs connect RNA-based gene regulation with chromatin-associated epigenetic mechanisms.

RNA-Directed DNA Methylation (RdDM) in Plants

RNA-directed DNA methylation (RdDM) is a plant epigenetic pathway in which small interfering RNAs guide DNA methylation machinery to specific genomic regions.

Main stages of canonical RdDM:

1. RNA Polymerase IV (Pol IV): Produces precursor transcripts from selected genomic regions.
2. RDR2: Converts these transcripts into double-stranded RNA.
3. DCL3: Processes double-stranded RNA into predominantly 24-nucleotide siRNAs.
4. AGO4/AGO6: Associates with siRNAs to form targeting complexes.
5. RNA Polymerase V (Pol V): Produces scaffold RNAs at target genomic regions.
6. DRM2: Catalyzes de novo cytosine methylation at targeted DNA sequences.

Biological significance:

- **Silencing of transposable elements and repetitive sequences.**
- **Establishment and maintenance of DNA methylation patterns.**
- **Regulation of selected genes and nearby regulatory regions.**
- **Contribution to genome stability and developmental regulation.**

Biological functions:

- **Regulation of developmental and stress-responsive genes.**
- **Control of transposable element activity.**
- **Recruitment or guidance of chromatin-modifying complexes in specific pathways.**
- **Maintenance of genome stability.**

Important distinction: Not all non-coding RNAs regulate chromatin directly; many function primarily through post-transcriptional mechanisms.

Key concept: Regulatory non-coding RNAs connect RNA-based gene regulation with chromatin-associated epigenetic mechanisms.

RNA-Directed DNA Methylation (RdDM) in Plants

RNA-directed DNA methylation (RdDM) is a plant epigenetic pathway in which small interfering RNAs guide DNA methylation machinery to specific genomic regions.

Main stages of canonical RdDM:

1. RNA Polymerase IV (Pol IV): Produces precursor transcripts from selected genomic regions.
2. RDR2: Converts these transcripts into double-stranded RNA.
3. DCL3: Processes double-stranded RNA into predominantly 24-nucleotide siRNAs.
4. AGO4/AGO6: Associates with siRNAs to form targeting complexes.
5. RNA Polymerase V (Pol V): Produces scaffold RNAs at target genomic regions.
6. DRM2: Catalyzes de novo cytosine methylation at targeted DNA sequences.

Biological significance:

- Silencing of transposable elements and repetitive sequences.
- Establishment and maintenance of DNA methylation patterns.
- Regulation of selected genes and nearby regulatory regions.
- Contribution to genome stability and developmental regulation.

RNA-Directed DNA Methylation (RdDM) in Plants

Simplified canonical RdDM pathway



Key concept: RdDM provides sequence-directed epigenetic regulation by connecting small RNAs with DNA methylation machinery.

ChIP-seq: Analysis of Histone Modifications and Protein–DNA Interactions

Chromatin immunoprecipitation followed by sequencing (ChIP-seq) is a high-throughput technique used to identify genomic regions associated with specific histone modifications or DNA-binding proteins.

Main experimental stages:

1. Chromatin preparation: Isolation and fragmentation of chromatin from biological samples.
2. Immunoprecipitation: Enrichment of chromatin fragments using antibodies against a selected histone modification or DNA-associated protein.
3. DNA purification: Recovery of DNA fragments associated with the target.
4. Library preparation and sequencing: High-throughput sequencing of enriched DNA.
5. Bioinformatics analysis: Read alignment, peak detection, and genomic annotation.

Major applications:

- Genome-wide mapping of histone modifications, including H3K4me3, H3K27me3, and H3K9me2.
- Identification of genomic regions occupied by transcription factors.
- Investigation of chromatin states during plant development and stress responses.
- Integration of chromatin profiles with RNA-seq data.

ChIP-seq workflow

Chromatin isolation and fragmentation

Antibody-based immunoprecipitation

DNA purification and sequencing

Read alignment and peak calling

Genome-wide protein–DNA association map

Key concept: ChIP-seq identifies genomic regions enriched for a particular chromatin-associated protein or histone modification.

ATAC-seq: Chromatin Accessibility Profiling

Assay for Transposase-Accessible Chromatin using sequencing (ATAC-seq) is a technique used to identify genomic regions with accessible chromatin.

Principle: A hyperactive Tn5 transposase inserts sequencing adapters preferentially into accessible DNA regions, enabling their identification by high-throughput sequencing.

Main experimental stages:

1. Isolation of intact nuclei from cells or plant tissues.
2. Tn5-mediated DNA fragmentation and adapter insertion (tagmentation).
3. Library amplification and high-throughput sequencing.
4. Read alignment and identification of accessible chromatin regions.
5. Functional annotation of regulatory DNA elements.

Major applications:

- Identification of accessible promoters and potential enhancers.
- Investigation of chromatin changes during development.
- Analysis of chromatin accessibility under environmental stress.
- Integration with transcriptomic data to prioritize regulatory regions.

Advantages:

- Relatively low sample input requirements.
- Efficient preparation compared with some traditional chromatin accessibility assays.
- Applicable to bulk and single-cell analyses.

Key concept: ATAC-seq reveals chromatin accessibility but does not directly establish transcription factor binding or gene activation.



Integration and Bioinformatics Analysis of Epigenomic Data

Epigenomic data integration combines information from DNA methylation, histone modifications, chromatin accessibility, and gene expression to investigate genome regulation.

Main analytical approaches:

- DNA methylation analysis: Identification of methylated cytosines and differentially methylated regions (DMRs).
- ChIP-seq analysis: Detection and annotation of genomic regions enriched for histone modifications or chromatin-associated proteins.
- ATAC-seq analysis: Identification of accessible chromatin regions and potential regulatory elements.
- RNA-seq integration: Comparison of epigenomic patterns with transcript abundance.
- Functional analysis: Gene Ontology, pathway enrichment, and regulatory network analysis.

Common bioinformatics tools:

- Bismark — bisulfite sequencing alignment and methylation calling.
- MACS3 — peak detection for ChIP-seq and ATAC-seq.
- deepTools — visualization and comparison of genomic signals.
- IGV — interactive visualization of sequencing data.

Example in plant biotechnology: Combining DNA methylation profiling, ATAC-seq, and RNA-seq can help identify candidate regulatory regions associated with drought-responsive gene expression in wheat

Integrated epigenomic analysis

DNA methylation **Histone modifications**

Chromatin accessibility **RNA expression**

Integrated bioinformatics analysis

Candidate regulatory mechanisms and functional interpretation

Key concept: Integrating multiple epigenomic datasets provides a more comprehensive understanding of genome regulation than analyzing individual molecular layers separately.

Epigenetic Regulation of Plant Growth and Development

Epigenetic regulation plays an essential role in coordinating gene expression during plant growth, differentiation, and developmental transitions.

Major developmental processes:

- Seed germination: Chromatin modifications and DNA methylation contribute to the regulation of dormancy and germination-associated genes.
- Root and shoot development: Epigenetic mechanisms influence cell differentiation, meristem activity, and organ formation.
- Flowering regulation: Histone modifications, particularly H3K27me3, contribute to the regulation of flowering-related genes.
- Reproductive development: DNA methylation and chromatin remodeling participate in gametogenesis, embryogenesis, and seed development.
- Cellular differentiation: Tissue-specific chromatin states help establish and maintain developmental gene expression patterns.

Example: In Arabidopsis thaliana, prolonged cold exposure (vernalization) promotes epigenetic silencing of the flowering repressor FLOWERING LOCUS C (FLC), involving H3K27me3 accumulation.



Key concept: Epigenetic mechanisms coordinate developmental gene expression and contribute to plant developmental plasticity.

Epigenetic Responses to Abiotic and Biotic Stress

Environmental stresses can induce changes in DNA methylation, histone modifications, and chromatin accessibility, influencing plant gene expression.

Major stress-related epigenetic responses:

- Drought stress: Changes in DNA methylation and histone marks may accompany the regulation of dehydration-responsive genes.
- Salinity stress: Chromatin remodeling and methylation changes can contribute to transcriptional adjustments associated with ion homeostasis.
- Temperature stress: Histone modifications participate in regulating heat- and cold-responsive genes.
- Pathogen infection: Epigenetic mechanisms can influence defense-associated transcriptional responses.
- Oxidative stress: Chromatin-associated regulation may contribute to changes in antioxidant and stress signaling pathways.

Potential regulatory targets:

- Stress-responsive transcription factors (DREB, NAC, WRKY).
- Genes involved in osmotic adjustment.
- Antioxidant defense-associated genes.
- Hormone signaling and defense pathways.

Key concept: Stress-associated epigenetic changes can influence transcriptional regulation, but their functional significance must be established experimentally.

Epigenetic Variation and Stress Memory

Epigenetic stress memory refers to the persistence of selected chromatin or DNA methylation states after an environmental stress has ended, potentially modifying responses to subsequent stress exposure.

Main characteristics:

- Some stress-induced epigenetic changes persist during recovery.
- Histone modifications and chromatin accessibility may contribute to transcriptional memory.
- DNA methylation changes can be associated with prolonged regulatory responses.
- Stress memory may be transient or maintained through mitotic cell divisions.
- The persistence and functional significance of memory depend on the species, tissue, and stress conditions.

Potential biological significance:

- Faster activation of selected stress-responsive genes.
- Modified responses to repeated environmental stress.
- Contribution to physiological acclimation and stress adaptation.

Key concept: Epigenetic stress memory may contribute to plant acclimation, but persistence of an epigenetic mark does not necessarily demonstrate improved stress tolerance.

Epigenetic Inheritance and Transgenerational Effects

Epigenetic inheritance is the transmission of certain regulatory states through cell divisions and, in some cases, across generations without changes in the underlying DNA sequence.

Major forms:

1. Mitotic inheritance: Maintenance of epigenetic states during somatic cell division.
2. Intergenerational effects: Effects observed in offspring that may result from direct parental or offspring exposure.
3. Transgenerational inheritance: Persistence of an epigenetic state in generations not directly exposed to the original environmental stimulus.

Potential mechanisms:

- Maintenance of DNA methylation patterns.
- Persistence or re-establishment of selected chromatin states.
- Small RNA-mediated epigenetic regulation.
- Biological significance:
- Maintenance of cellular identity.
- Developmental regulation.
- Formation of stable epialleles.
- Potential contribution to phenotypic diversity.

Important limitations:

- Many stress-induced epigenetic changes are not stably inherited.
- Genetic variation and parental environmental effects must be distinguished from epigenetic inheritance.
- Multigenerational experiments are necessary to demonstrate transgenerational effects.

Example: Naturally occurring and experimentally induced epialleles in plants can produce heritable differences in gene expression and phenotype.

Key concept: Epigenetic inheritance can contribute to phenotypic variation, but stable transgenerational transmission is not a universal property of stress-induced epigenetic changes.

Applications of Epigenomics in Plant Biotechnology

Epigenomics provides new opportunities for understanding and improving agriculturally important plant traits through the regulation of genome activity.

Major applications:

- Crop improvement: Identification of epigenetic variation associated with yield, grain quality, and developmental characteristics.
- Abiotic stress tolerance: Investigation of epigenetic mechanisms associated with drought, salinity, and temperature responses.
- Disease resistance: Characterization of chromatin modifications and DNA methylation patterns associated with plant defense.
- Epigenetic breeding: Exploration of stable epialleles and epigenetic diversity as complementary resources for crop improvement.
- Plant tissue culture: Analysis of epigenetic changes during regeneration, somatic embryogenesis, and micropropagation.
- Functional genomics: Identification of regulatory elements influencing agronomically important genes.

- Example in wheat: Integrating DNA methylation profiling with RNA-seq can identify candidate epigenetic regulatory mechanisms associated with drought-responsive gene expression.

Key concept: Epigenomics complements conventional breeding and molecular biotechnology by revealing regulatory variation beyond DNA sequence polymorphisms.

Applications in Medical and Industrial Biotechnology

Epigenomic technologies are increasingly applied in medicine, pharmaceutical research, and industrial biotechnology.

Medical biotechnology:

- Cancer research: Identification of abnormal DNA methylation and histone modification patterns.
- Disease biomarkers: Development of epigenetic markers for disease classification, diagnosis, and prognosis.
- Drug development: Investigation of DNA methyltransferase and histone deacetylase inhibitors.
- Precision medicine: Integration of epigenomic profiles with genomic and transcriptomic information.

Industrial biotechnology:

- Investigation of chromatin-mediated regulation in industrial fungi and other eukaryotic production systems.
- Regulation of secondary metabolite biosynthesis.
- Optimization of gene expression in selected microbial production strains.
- Identification of epigenetic factors affecting cellular differentiation and productivity.

Key concept: Epigenomic approaches support the identification of regulatory mechanisms and the development of new diagnostic and biotechnological strategies.

Challenges and Future Perspectives: Epigenome Editing

Epigenome editing involves targeted modification of epigenetic states at selected genomic regions without necessarily changing the underlying DNA sequence.

Emerging technologies:

- CRISPR/dCas9-based epigenome editing: Targeted recruitment of epigenetic regulatory proteins.
- Targeted DNA methylation: Experimental modification of methylation patterns at specific genomic loci.
- Targeted histone modification: Recruitment of histone-modifying enzymes to regulatory regions.
- Single-cell epigenomics: Investigation of cell-specific epigenetic states.
- Multi-omics integration: Combining epigenomic, transcriptomic, and phenotypic information.

Major challenges:

- Distinguishing causal epigenetic changes from correlated changes.
- Achieving high targeting specificity.
- Understanding the stability and inheritance of edited epigenetic states.
- Validating functional effects across developmental stages and environments.
- Improving delivery and regeneration systems in crop plants.

Future perspective: Integration of epigenome editing, artificial intelligence, and high-throughput phenotyping may facilitate the discovery and functional validation of regulatory mechanisms relevant to crop improvement.

Key concept: Epigenome editing offers a promising approach for studying and potentially modifying gene regulation, although many agricultural applications remain experimental.

Conclusion

- Epigenomics investigates genome-wide regulatory mechanisms that influence gene expression without necessarily altering DNA sequences.
- DNA methylation, histone modifications, chromatin remodeling, and regulatory RNAs contribute to the control of genome activity.
- ChIP-seq, ATAC-seq, and DNA methylation sequencing enable comprehensive characterization of epigenetic landscapes.
- Epigenetic mechanisms participate in plant development, environmental responses, and selected forms of cellular memory and inheritance.
- Integration of epigenomics with transcriptomics and other omics technologies supports functional genomics and biotechnology.
- Targeted epigenome editing represents an emerging strategy for investigating gene regulation and developing novel biotechnological approaches.

Overall conclusion: Epigenomics provides an essential framework for understanding dynamic genome regulation and expands the possibilities of postgenomic research.

Control Questions

1. What are epigenetics and epigenomics, and how do they differ from classical genetics?
2. What are the major mechanisms of epigenetic regulation?
3. How do euchromatin and heterochromatin differ in structure and function?
4. What are the main DNA methylation contexts in plants?
5. How do histone acetylation and methylation influence gene expression?
6. What is the role of non-coding RNAs in epigenetic regulation?
7. What are the major stages of RNA-directed DNA methylation?
8. What are the principles and applications of ChIP-seq and ATAC-seq?
9. How do epigenetic mechanisms contribute to plant development and stress responses?
10. What are the opportunities and limitations of epigenome editing in biotechnology?

Recommended Literature

- 1. Zhang, H., Lang, Z., & Zhu, J.-K. (2018). Dynamics and function of DNA methylation in plants. Nature Reviews Molecular Cell Biology, 19, 489–506. https://doi.org/10.1038/s41580-018-0016-z
- 2. Matzke, M. A., & Mosher, R. A. (2014). RNA-directed DNA methylation: An epigenetic pathway of increasing complexity. Nature Reviews Genetics, 15, 394–408. https://doi.org/10.1038/nrg3683
- 3. Law, J. A., & Jacobsen, S. E. (2010). Establishing, maintaining and modifying DNA methylation patterns in plants and animals. Nature Reviews Genetics, 11, 204–220. https://doi.org/10.1038/nrg2719
- 4. Buenrostro, J. D., et al. (2013). Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. Nature Methods, 10, 1213–1218. https://doi.org/10.1038/nmeth.2688
- 5. Springer, N. M., & Schmitz, R. J. (2017). Exploiting induced and natural epigenetic variation for crop improvement. Nature Reviews Genetics, 18, 563–575. https://doi.org/10.1038/nrg.2017.45
- 6. Gallego-Bartolomé, J. (2020). DNA methylation in plants: Mechanisms and tools for targeted manipulation. New Phytologist, 227, 38–44. https://doi.org/10.1111/nph.16529