Lecture 9. Phenomics and High-Throughput Phenotyping Technologies

Phenome, phenotyping platforms, RGB, fluorescence, thermal and hyperspectral imaging, sensors, digital phenotyping, plant phenomics, and genotype–phenotype relationships.

1. Aim of the lecture

- To introduce the principles of phenomics and high-throughput phenotyping, explain modern imaging and sensor-based technologies for quantitative assessment of biological traits, and demonstrate their applications in investigating genotype–phenotype relationships, plant stress responses, and crop improvement.

2. Expected learning outcomes

1. Define the phenome, phenotype, and phenomics.
2. Explain the relationship between genotype, environment, and phenotype.
3. Distinguish traditional and high-throughput phenotyping.
4. Describe greenhouse, laboratory, and field phenotyping platforms.
5. Explain RGB, chlorophyll fluorescence, thermal, multispectral, and hyperspectral imaging.
6. Identify the major types of sensors used in plant phenotyping.
7. Explain digital image processing and trait extraction.
8. Describe applications of machine learning and artificial intelligence in phenomics.
9. Discuss how phenotyping can be integrated with genomics, transcriptomics, proteomics, and metabolomics.
10. Explain the role of phenomics in crop breeding and precision biotechnology.

Lecture plan

- Introduction to phenome and phenomics
- What is the phenome?
- The phenome is the complete set of observable or measurable characteristics of an organism, resulting from the interaction of its genetic makeup, environmental conditions, and developmental processes.
- The phenome includes:
  - Morphological traits: plant height, leaf area, grain size.
  - Physiological traits: photosynthesis, transpiration, water status.
  - Biochemical traits: pigment content, enzyme activities, metabolites.
  - Developmental traits: flowering time, maturation.
  - Agronomic traits: biomass, yield, grain quality.
- Key concept: The phenome represents the measurable manifestation of biological processes at multiple levels.

Introduction to phenome and phenomics

What is the phenome?

The phenome is the complete set of observable or measurable characteristics of an organism, resulting from the interaction of its genetic makeup, environmental conditions, and developmental processes.

The phenome includes:

1. Morphological traits: plant height, leaf area, grain size.
2. Physiological traits: photosynthesis, transpiration, water status.
3. Biochemical traits: pigment content, enzyme activities, metabolites.
4. Developmental traits: flowering time, maturation.
5. Agronomic traits: biomass, yield, grain quality.

Key concept: The phenome represents the measurable manifestation of biological processes at multiple levels.

What is phenomics?

Phenomics is the comprehensive, systematic, and often high-throughput measurement and analysis of phenotypic traits across organisms, developmental stages, and environmental conditions.

Its main objectives are to:

1. *Quantify phenotypic diversity.*
2. *Characterize growth and development.*
3. *Identify responses to environmental factors.*
4. *Investigate genotype–phenotype relationships.*
5. *Discover phenotypic biomarkers.*
6. *Support genetic research and breeding.*

Phenotype as a product of genotype and environment

The phenotype is influenced by genetic, environmental, developmental, and regulatory factors.

A simplified quantitative-genetic model is:

$$P = G + E + G \times E + e$$

- Where:
- P — observed phenotypic value.
- G — genetic contribution.
- E — environmental contribution.
- (G x E) — genotype-by-environment interaction.
- *E* — residual variation.

This is a statistical decomposition of phenotypic variation, not a universal molecular equation.

Position of phenomics in postgenomic biology

- **Genome — genetic potential**
- **Transcriptome — gene expression**
- **Proteome — functional proteins**
- **Metabolome — biochemical processes**
- **Phenome — measurable traits**
- **Environment, development and regulatory feedback influence every level.**

Major objectives of plant phenomics

Plant phenomics aims to:

1. *Measure growth and development objectively.*
2. *Quantify physiological and morphological variation.*
3. *Identify stress-responsive traits.*
4. *Analyze phenotypic plasticity and stability.*
5. *Link genomic variation to measurable traits.*
6. *Improve selection efficiency in plant breeding.*

2. Traditional and high-throughput phenotyping. Traditional phenotyping methods

Traditional plant phenotyping commonly includes manual or instrument-assisted measurements of:

1. Plant height and tiller number.
2. Leaf area and biomass.
3. Chlorophyll content.
4. Relative water content.
5. Flowering and maturity dates.
6. Grain number, size, and weight.
7. Disease severity.

These measurements remain important reference methods for validating automated approaches.

High-throughput phenotyping (HTP)

High-throughput phenotyping refers to automated or semi-automated measurement of many phenotypic traits across large numbers of samples.

Advantages include:

- Increased measurement capacity.
- Reduced manual labor.
- Repeated measurements over time.
- Quantitative digital records.
- Potentially improved consistency.
- Integration with genomic and environmental datasets.

Main phenotyping platforms

Platform	Main applications
Laboratory platforms	Seeds, roots, leaves, cellular traits
Greenhouse platforms	Controlled growth and stress experiments
Field platforms	Agronomic traits under realistic conditions
UAV-based platforms	Canopy monitoring over large field areas
Ground-based robotic platforms	Close-range crop measurements

Controlled-environment versus field phenotyping

Controlled environments

- More precise regulation of temperature, light, and water.
- Easier repeated imaging.
- Reduced environmental variability.
- Limited representation of field conditions.

Field environments

- More realistic genotype × environment interactions.
- Larger spatial and temporal variability.
- Challenges from wind, sunlight, and background interference.
- Greater relevance for breeding performance.

3. Imaging technologies in plant phenotyping: RGB and fluorescence

Imaging technologies enable non-destructive assessment of plant morphology, physiology, and biochemical characteristics.

- The principal methods include:
- RGB,
- chlorophyll fluorescence,
- thermal,
- multispectral, and
- hyperspectral imaging.

RGB imaging

RGB imaging uses three visible-light channels: red, green, and blue.

It is used to measure:

- Projected leaf area and canopy cover.
- Plant height and architecture.
- Growth dynamics.
- Leaf color and senescence.
- Grain size and shape.

RGB imaging is relatively inexpensive and suitable for automated analysis, although estimates of biomass and other traits require appropriate calibration.

Chlorophyll fluorescence imaging

Chlorophyll fluorescence imaging evaluates the functioning of the photosynthetic apparatus, particularly photosystem II (PSII).

Important parameters include:

- Fv/Fm: maximum quantum efficiency of PSII in dark-adapted samples.
- Y(II): effective quantum yield of PSII.
- NPQ: non-photochemical quenching.
- ETR: estimated electron transport rate.

Applications include detection of photosynthetic stress, drought responses, photoinhibition, and disease-associated changes.

Fluorescence imaging in stress research

Fluorescence imaging can reveal spatial variation in photosynthetic performance before severe visible symptoms appear.

- Stress treatment → changes in PSII activity → fluorescence response → spatial visualization → physiological interpretation

Fluorescence measurements are especially useful when combined with water status, pigment content, and growth measurements.

4. Thermal, multispectral and hyperspectral imaging

Thermal imaging imaging measures infrared radiation emitted by plant surfaces and estimates their temperature.

Leaf temperature is influenced by transpiration, stomatal conductance, radiation, and environmental conditions.

Applications:

- Drought and heat stress detection.
- Canopy temperature assessment.
- Screening for water-use-related traits.
- Identification of spatial temperature variation.

Thermal imaging and plant water status

Under comparable environmental conditions, reduced transpiration may increase leaf temperature.

- Water deficit → stomatal closure → reduced evaporative cooling → increased leaf temperature
- However, temperature alone does not provide a direct measurement of stomatal conductance or drought tolerance. Environmental conditions and canopy structure must also be considered.

Multispectral imaging

Multispectral imaging measures reflectance in a limited number of selected wavelength bands, commonly including visible and near-infrared regions.

It is used to estimate vegetation indices and investigate:

- Canopy greenness.
- Pigment-related traits.
- Vegetation cover.
- Stress-associated spectral changes.
- A common vegetation index is:

NDVI=NIR-Red/NIR+Red

where NIR represents near-infrared reflectance.

Hyperspectral imaging

- Hyperspectral imaging records reflectance across numerous narrow, often contiguous wavelength bands.
- It produces a three-dimensional dataset:
- X position × Y position × wavelength
- Applications include:
- Pigment estimation.
- Detection of biochemical and structural variation.
- Stress and disease classification.
- Investigation of plant water and nutritional status.

Comparison of imaging methods

Method	Main information	Typical application
RGB	Visible morphology and color	Growth, architecture, grain shape
Fluorescence	Photosynthetic performance	PSII efficiency, stress
Thermal	Surface temperature	Water stress, canopy cooling
Multispectral	Reflectance in selected bands	Vegetation indices
Hyperspectral	Detailed spectral signatures	Biochemical and stress-related traits

5. Sensors and digital phenotyping

Sensors in plant phenotyping measure plant properties and environmental conditions.

Major groups include:

1. **Optical sensors.**
2. **Temperature and humidity sensors.**
3. **Soil moisture sensors.**
4. **Gas-exchange instruments.**
5. **Chlorophyll and fluorescence sensors.**
6. **LiDAR and depth sensors.**

Digital phenotyping

Digital phenotyping uses imaging, sensors, automated data collection, and computational analysis to quantify biological traits.

Its general workflow is:

- **Plant material**
- **Imaging and sensor measurements**
- **Image processing and segmentation**
- **Quantitative trait extraction**
- **Statistical analysis and interpretation**

Image processing and trait extraction

Typical image-analysis operations include:

- Image calibration.
- Background removal.
- Plant segmentation.
- Object identification.
- Measurement of geometric characteristics.
- Extraction of color, texture, and spectral features.

These methods transform images into numerical phenotypic datasets.

High-Throughput Phenotyping in Microorganisms

High-throughput phenotyping (HTP) in microorganisms enables rapid, automated characterization of microbial growth, metabolism, stress tolerance, and functional traits across large numbers of strains or experimental conditions.

Major technologies:

Automated microplate readers: Measure microbial growth kinetics, optical density, fluorescence, and metabolic activity.

- Phenotype microarrays: Characterize substrate utilization, nutrient requirements, and responses to chemical compounds.
- Flow cytometry: Analyzes cell size, fluorescence, viability indicators, and population heterogeneity.
- Microfluidic platforms: Enable single-cell monitoring of microbial growth and responses to environmental changes.
- Automated colony imaging: Quantifies colony size, morphology, pigmentation, and growth characteristics.
- High-content microscopy: Provides quantitative information on microbial cell morphology and cellular organization.

- Major applications:
- Screening microbial strains for industrial biotechnology.
- Identification of stress-tolerant microorganisms.
- Optimization of fermentation processes.
- Characterization of metabolic diversity and substrate utilization.
- Functional analysis of mutant libraries and gene-edited strains.

Example: Automated phenotyping of Saccharomyces cerevisiae strains can identify differences in growth rates, fermentation performance, and tolerance to ethanol or osmotic stress.

Key concept: Microbial HTP combines automation, miniaturization, and quantitative analysis to accelerate functional genomics and microbial strain improvement.

High-Throughput Phenotyping in Animals and Other Model Organisms

High-throughput phenotyping in animals and model organisms integrates automated imaging, sensor technologies, and computational analysis to characterize physiological, developmental, and behavioral traits.

Major technologies:

- Automated imaging systems: Monitor morphology, growth, development, and movement.
- High-content microscopy: Quantifies cellular phenotypes in cultured cells and small model organisms.
- Behavioral tracking systems: Analyze locomotion, activity patterns, and responses to stimuli.
- Wearable and environmental sensors: Measure physiological and behavioral parameters.
- Microfluidic screening: Enables automated phenotyping of small organisms and cellular models.
- AI-assisted image analysis: Supports automated classification and quantitative phenotype extraction.

Major model organisms and applications

Organism	Major phenotypic traits
Caenorhabditis elegans	Development, movement, lifespan, stress responses
Drosophila melanogaster	Development, behavior, morphology
Danio rerio (zebrafish)	Embryonic development, organ formation, behavior
Laboratory mice	Metabolism, physiology, behavior, disease phenotypes

Artificial intelligence in phenomics

Machine learning and deep learning can assist in:

- Plant and organ segmentation.
- Disease symptom recognition.
- Grain counting and classification.
- Stress detection.
- Trait prediction.
- Analysis of large imaging datasets.

Models must be validated across relevant genotypes, environments, and imaging conditions to assess their reliability.

6. Plant phenomics and genotype–phenotype relationships

Plant phenomics integrates high-throughput measurements of morphological, physiological, and agronomic traits.

It provides quantitative phenotypes for:

- **GWAS and QTL mapping.**
- **Functional genomics.**
- **Genomic selection.**
- **Mutation breeding.**
- **Gene-editing studies.**

Genotype–phenotype relationships

- Genetic variation
- Gene expression and molecular regulation
- Protein and metabolic processes
- Physiological and developmental traits
- Phenotype
- Environmental conditions influence all stages.

Phenomics and genotype × environment interaction

- High-throughput phenotyping allows researchers to compare genotypes across environments and developmental stages.

Repeated measurements can reveal:

- Growth trajectories.
- Stress-response dynamics.
- Phenotypic plasticity.
- Trait stability.
- Genotype-specific environmental responses.

Applications and challenges

Applications in crop improvement

- Phenomics contributes to:
- Identification of superior genotypes.
- Screening for abiotic and biotic stress responses.
- Evaluation of mutant populations.
- Identification of marker–trait associations.
- Selection for yield and grain quality.
- Integration of phenotypic and genomic datasets.

. Challenges and future perspectives

Major challenges include:

- Environmental variability.
- Differences in sensor calibration.
- Large data volumes.
- Image segmentation errors.
- Standardization of phenotyping protocols.
- Transferability of models across environments.

Future directions include multimodal sensing, robotic platforms, AI-assisted phenotyping, and integration with multi-omics datasets.

Challenges and Future Perspectives in Single-Cell and Spatial Omics

Single-cell and spatial omics technologies are transforming postgenomic research by enabling high-resolution investigation of cellular heterogeneity, tissue organization, and molecular regulation.

Major challenges:

- Sample preparation: Difficulties in isolating viable cells or intact nuclei, particularly from plant tissues with rigid cell walls.
- Technical variability: Low RNA capture efficiency, dropout events, doublets, and batch effects.
- Spatial resolution: Trade-offs between molecular coverage, sensitivity, and spatial precision.
- Data complexity: Large datasets requiring advanced bioinformatics and computational resources.
- Biological interpretation: Challenges in identifying cell types, reconstructing developmental trajectories, and validating cell–cell interactions.

Future research directions:

- Single-cell multi-omics: Simultaneous profiling of transcriptomic, epigenomic, and selected proteomic features.
- High-resolution spatial transcriptomics: Improved mapping of gene expression at cellular and subcellular resolution.
- Spatial proteomics and metabolomics: Investigation of protein and metabolite distributions within tissues.
- AI-assisted analysis: Automated cell annotation, spatial pattern recognition, and multimodal data integration.
- Plant biotechnology: Development of cell-type-specific molecular maps to support crop improvement and stress research.

Key concept: Integration of single-cell, spatial, and multi-omics technologies will enable increasingly comprehensive analysis of biological processes at cellular and tissue levels.

Conclusion

- Single-cell omics reveals molecular heterogeneity among individual cells and nuclei.
- Spatial omics maps molecular features within their original tissue context.
- Droplet-based sequencing enables high-throughput characterization of cellular transcriptomes.
- Single-cell epigenomics and proteomics provide complementary information about chromatin regulation and protein expression.
- Spatial transcriptomics, proteomics, and metabolomics reveal tissue-specific molecular organization.
- Bioinformatics methods support cell clustering, cell-type identification, developmental trajectory inference, and spatial data analysis.
- Integration of single-cell and spatial omics advances understanding of development, environmental responses, and cellular interactions.

Overall conclusion: Single-cell and spatial omics connect molecular regulation with cellular identity and tissue organization, providing powerful approaches for functional genomics, systems biology, and modern biotechnology.

Control Questions

1. What are single-cell and spatial omics, and how do they differ from bulk omics?
2. What are the major advantages of single-cell RNA sequencing?
3. How does droplet-based single-cell sequencing work?
4. What are cell barcodes and unique molecular identifiers (UMIs)?
5. What are the main applications of single-cell epigenomics and proteomics?
6. How do scATAC-seq and single-cell RNA-seq provide complementary information?
7. What are the major principles and technologies of spatial transcriptomics?
8. How are cell clustering, dimensionality reduction, and trajectory analysis applied to single-cell data?
9. What challenges are associated with single-cell analysis of plant tissues?
10. How can single-cell and spatial omics contribute to plant biotechnology and crop improvement?

Recommended Literature

- 1. Stuart, T., & Satija, R. (2019). Integrative single-cell analysis. Nature Reviews Genetics, 20, 257–272. https://doi.org/10.1038/s41576-019-0093-7
- 2. Luecken, M. D., & Theis, F. J. (2019). Current best practices in single-cell RNA-seq analysis: A tutorial. Molecular Systems Biology, 15, e8746. https://doi.org/10.15252/msb.20188746
- 3. Marx, V. (2021). Method of the Year: Spatially resolved transcriptomics. Nature Methods, 18, 9–14. https://doi.org/10.1038/s41592-020-01033-y
- 4. Longo, S. K., Guo, M. G., Ji, A. L., & Khavari, P. A. (2021). Integrating single-cell and spatial transcriptomics to elucidate intercellular tissue dynamics. Nature Reviews Genetics, 22, 627–644. https://doi.org/10.1038/s41576-021-00370-8
- 5. Denyer, T., et al. (2019). Spatiotemporal developmental trajectories in the Arabidopsis root revealed using high-throughput single-cell RNA sequencing. Developmental Cell, 48, 840–852.e5. https://doi.org/10.1016/j.devcel.2019.02.022
- 6. Ryu, K. H., et al. (2019). Single-cell RNA sequencing resolves molecular relationships among individual plant cells. Plant Physiology, 179, 1444–1456. https://doi.org/10.1104/pp.18.01482
- 7. Shaw, R., Tian, X., & Xu, J. (2021). Single-cell transcriptome analysis in plants: Advances and challenges. Molecular Plant, 14, 115–126. https://doi.org/10.1016/j.molp.2020.10.012